



ORIGINAL ARTICLE

Improving Diabetic Neuropathy Knowledge in an Urban Elderly Population: A Community-Based Intervention Study

Laurentius Johan Ardian^{1*}, Aktaruddin Arief Santoso¹, Hilmia Fahma¹, Anoraga Mona Cinta Gadis Cantika Gusyono¹, Satria Fudi Verbimanto¹

¹Faculty of Medicine Universitas Pembangunan Nasional Veteran Jawa Timur, Surabaya, Indonesia

ARTICLE INFO

Correspondence:

Laurentius Johan Ardian
Faculty of Medicine Universitas
Pembangunan Nasional Veteran Jawa
Timur, Surabaya, Indonesia

Email:

laurentius.johan.ardian.fk@upnjatim.ac.id

Article history:

Received : August 29, 2025

Received in revised form :

October 13, 2025

Accepted : October 15, 2025

Keywords:

Diabetes
Elderly
Health education

ABSTRACT

Background : The global population is aging rapidly, with older adults increasingly affected by chronic conditions such as diabetes mellitus. In Indonesia, diabetes among the elderly poses a significant public health concern, often complicated by under-recognized conditions like diabetic neuropathy. **Aim.** This study aimed to assess the effectiveness of a community-based intervention focused on targeted health education on diabetic neuropathy among elderly residents in urban area.

Methods : A pre- and post-test design was used to assess the intervention's effectiveness. In collaboration with local health cadres and facilities, participants completed identical questionnaires on diabetic neuropathy before and after the health education session to measure improvements in knowledge.

Results : A total of 143 elderly participants (median age: 65 years) were enrolled, with 73.4% being female. Post-education test scores showed a statistically significant improvement compared to pre-test scores ($p < 0.001$, Wilcoxon test). Of the participants, 22.4% were newly identified as having diabetes, and another 22% fell into the prediabetes range. Despite some variations in learning outcomes, 84.6% showed improved scores, indicating high effectiveness of the health education session.

Conclusion : This study demonstrates that a community-based approach combining early diabetes screening and health education significantly improves elderly participants' knowledge about diabetes and diabetic neuropathy.

Medical and Health Science Journal

<https://doi.org/10.33086/mhsj.v9i2.7974>



This is an open access article distributed under the Creative Commons Attribution-ShareAlike 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. ©2025The Author(s)

Introduction

The world is experiencing unprecedented demographic aging, with the proportion of individuals aged ≥ 60 years increasing at an accelerated rate. By 2020, the global population of older adults surpassed the number of children under five years of age.¹ Projections indicate that between 2015 and 2050, the proportion of the world's elderly population will nearly double, rising from 12% to 22%.² Indonesia exemplifies this trend, ranking among the top eight countries with the largest elderly population globally. With 8.2% of its population aged 60 years or older in 2015, Indonesia has officially transitioned into an aging society.³

Aging is associated with a progressive decline in physiological resilience, leading to increased susceptibility to chronic diseases and organ dysfunctions. Immunosenescence—the age-related weakening of the immune system—increases vulnerability to infections, malignancies, and inflammatory conditions, particularly cardiovascular diseases.^{4,5} Furthermore, degenerative disorders such as Alzheimer's disease, Parkinson's disease, and osteoarthritis are characterized by chronic inflammation that exacerbates tissue damage.⁶ These age-related changes underscore the need for targeted healthcare strategies to mitigate morbidity in older populations.

Diabetes mellitus is a major health concern among the elderly in Indonesia. According to Riskesdas 2018, diabetes prevalence in individuals aged 55–64, 65–74, and ≥ 75 years was 6.3%, 6.0%, and 3.0%, respectively.⁷ Moreover, diabetes can lead to serious complications such as diabetic neuropathy. Although common among people with diabetes, diabetic neuropathy is often under-recognized and poorly understood by both patients and caregivers. This lack of awareness can delay diagnosis and management, ultimately worsening the patient's quality of life.

The government has launched several initiatives, such as *Gerakan Masyarakat Hidup Sehat* (Germas); however, public awareness of diabetes as a non-communicable disease and its complications remains inadequate. Studies indicate that knowledge levels among urban diabetics are predominantly low, with limited formal education identified as a key contributing factor.^{8,9} This knowledge gap persists even in urban areas with greater access to health care.¹⁰

There is no doubt that diabetes education is essential for establishing foundational knowledge about diabetes management, both at the time of diagnosis and throughout the patient's journey as treatment needs and preferences evolve.¹¹ Evidence demonstrates that such education significantly

improves glycemic control, as measured by reductions in HbA1c levels.¹² However, the brevity of clinical encounters often restricts healthcare providers to superficial interventions, such as medication adjustments, glucose monitoring guidance, and behavioral recommendations, tasks hindered by varying levels of patient health literacy. Sustainable lifestyle modifications and community-based health promotion require extended support beyond clinical settings, necessitating collaborative programs tailored to high-risk and marginalized groups.¹³

Faculty of Medicine at Universitas Pembangunan Nasional "Veteran" Jawa Timur launched a health education program focused on diabetes education for elderly residents in Gunung Anyar District. Such interventions are vital for bridging the gap between clinical care and sustainable health outcomes in the aging population.

Methods

2.1 Study Design and Participants

Older adults aged 60 years or older who were present at the Gunung Anyar District Office on June 15th, 2025, were recruited for this study. A pre-test/post-test design was implemented. This one-group experimental approach involves taking measurements before and after an intervention to assess changes in knowledge level. We excluded individuals who could not complete both the pre- and post-test sessions, as well as those with severe cognitive impairment (defined as an Abbreviated Mental Test score ≤ 3). The sample size was calculated based on the mean difference and deviation standard between pre- and post-test values based on previous study.¹⁴ The estimated minimum sample size was 83 subjects. A two-way hypothesis test was employed, with alpha and beta values set at 5% and 10%, respectively. Ethical approval was obtained from the Research Ethics Committee at the Faculty of Medicine, Universitas Pembangunan Nasional "Veteran" Jawa Timur.

2.2 Baseline Data and Follow-up

Baseline data were collected using questionnaires at the start of the recruitment process. Data obtained from history-taking included: (i) demographic data (sex, age, educational background); (ii) diabetes status, categorized as diabetes, prediabetes, or normal; (iii) hypertension status, categorized as hypertensive or normal; (iv) cognitive function, assessed using the Abbreviated Mental Test (AMT) and classified as cognitive impairment (score 0–6) or normal (score 7–10); (v) polypharmacy, defined as the daily use

of five or more medications; and (vi) pre-test and post-test scores, obtained from a self-reported questionnaire completed by the participants.

2.3 Statistical Analysis

Data were analyzed using SPSS software version 24.0 (IBM, Armonk, New York, USA). First, a descriptive analysis of all variables was performed, including demographics and pre- and post-test scores. The paired t-test was used to compare scores if the data were normally distributed; otherwise, the Wilcoxon signed-rank test was applied. A further analysis was conducted to determine the proportion of participants who showed an improvement in their post-test score.

Results

3.1 Characteristics of the subjects

A total of 143 participants from the Gunung Anyar District attended the program. The median age of the participants was 65 (60-83) and most of the participants were female (73,4%). (Table 1) Most participants were not previously diagnosed with diabetes.

Table 1. Characteristics of The Subjects

Characteristics	n, %
Gender	
Male	38 (26,6)
Female	105 (73,4)
Age (Median, min-max)	65 (60-83)
Marital Status	
Married	114 (79,7)
Divorce	8 (5,6)
Divorced by death	21 (14,7)
Education	
Elementary school	12 (8,4)
Junior high school	42 (29,4)
Senior high school	69 (48,3)
Undergraduate above	20 (14)
Cognitive status	
Normal	128 (86,48)
Mild Cognitive Impairment	20 (13,5)
Random blood glucose (Median, min-max), mg/dL	138 (81-565)
Hypertension	
Yes	68 (45,9)
No	80 (54,05)
Diabetes mellitus status	
Diabetes	32 (22,4)
Pre-DM	31 (21,7)
Normal	80 (55,9)
Polypharmacy (>5 drugs)	
Yes	20 (13,79)
No	128 (86,4)
Using Diabetes Medication	
Yes	4 (2,8)
No	139 (97,2)

3.2. Knowledge Level Before and After Health Promotion and Education

Based on Table 2, there was a significant increase in knowledge level after health promotion and education ($p < 0,001$ Wilcoxon test).

Table 2. Score before and after Health Counselling

Pre-test score	Post-test score	P value
70 (0-100)	90 (40-100)	0,000*

*Wilcoxon test

Further analysis of the Wilcoxon test (Table 3) revealed that 121 (84,6%) subjects had an increased knowledge score, with a mean rank after health counselling 61,79. The mean rank before health counselling was 27.

Table 3. Further Analysis of Wilcoxon Test

	N	Mean Rank
Negative Ranks	1 ^a	27.00
Positive Ranks	121 ^b	61.79
Ties	21 ^c	
Total	143	

^a post-test score < pre-test score

^b post-test score > pre-test score

^c post-test score = pre-test score

Discussion

With the rapid growth of information accessibility today, people can obtain knowledge through various means. However, acquiring health information directly from healthcare facilities offers distinct advantages. Unlike unverified sources, guidance from medical professionals is evidence-based and tailored to individual needs. Additionally, health education empowers patients, improving their health-seeking behaviors¹⁵. Counseling further enhances this process by addressing specific concerns, enabling clients to resolve their issues effectively¹⁶.

The present study showed a statistically significant improvement in participants' knowledge. This was evidenced by a notable increase in post-test scores, as analysed using the Wilcoxon test ($p < 0.05$). These findings indicate that the educational intervention successfully enhanced the community's understanding of diabetes complications, particularly diabetic neuropathy. This result aligns with previous studies demonstrating that community-based health education can effectively increase health literacy and awareness of chronic complications of diabetes.¹⁷

Subsequent analysis indicated that one participant demonstrated a decline in performance, with a lower post-test score compared to the pre-test, while 21 participants exhibited no change in their test scores. These findings may be attributed to several factors, including limited motivation to engage with the educational material, low levels of formal education that may have impeded comprehension, or the use of health counselling methods that were not optimally aligned with

individual learning needs. It is well recognized that different educational approaches may be more effective for specific groups, depending on factors such as age, gender, educational background, and occupation.¹⁸

Our community development program served as an example of a Community-Based Health Intervention (CBHI). The participation of community-dwelling older adults in this initiative demonstrated promising results, indicating that the elderly exhibit proactive health-seeking behaviors. Their involvement in such programs was significantly associated with higher odds of undergoing regular blood pressure monitoring (adjusted odds ratio [OR], 3.09; 95% confidence interval [CI], 2.67–3.58) and blood glucose testing (adjusted OR, 1.88; 95% CI, 1.58–2.23)¹⁹. From a community development perspective, the active engagement of local health cadres and community members plays a critical role in strengthening local capacity for managing chronic diseases.²⁰

However, this study has limitations in terms of follow-up duration and did not assess long-term behavioral changes. Knowledge alone is insufficient to improve health behavior. The six steps of behavioral change are pre-contemplation, contemplation, preparation, action, maintenance, and termination. Knowledge from health counselling helps individuals recognize the need for change, helps individuals recognize the need for change, which is essential for moving from precontemplation (not considering change) to contemplation (thinking about change). (Fang, n.d.) Moreover, changing behavior itself depends not only on a patient's individual capacity to regulate impulses but also on ecosystems, including shared structural affordances and impediments, social norms, culture-wide practices, and more. Further research is needed to evaluate the sustained impact of health counselling on community practices and the incidence of diabetic neuropathy over time.²²

Health counselling faces several challenges. In general, there are two obstacles from the perspective of patients with diabetes regarding educational training. First, patients sometimes found it difficult to understand the material given by the healthcare team, and the topic of the counselling was boring and repeated. In addition to boring material, patients are sometimes afraid to ask. This problem also becomes an obstacle to receiving materials delivered during education. Thus, two-way interactive health counselling is needed, and the health counselling should consider patients' prior knowledge.²³

Overall, this study highlights that early detection and targeted health counselling on diabetic neuropathy are effective strategies to

improve public awareness and community readiness in managing one of the most debilitating complications of diabetes.

Conclusion

Our intervention effectively enhanced participants' knowledge of diabetes and its associated neurological complications. These short-term outcomes underscore the importance of collaborative efforts in promoting diabetes awareness and facilitating early diagnosis among elderly populations. To evaluate the long-term impact, future studies should include structured follow-up assessments to measure sustained knowledge retention, compliance with preventive practices, and potential declines in the incidence of diabetic neuropathy.

Conflict of Interest

No conflict of interest in this study

Acknowledgement

The authors gratefully acknowledge all participants for their invaluable contributions to this study. We extend our sincere appreciation to the Lembaga Penelitian dan Pengabdian Masyarakat (LPPM) of Universitas Pembangunan Nasional "Veteran" Jawa Timur for their financial support through the 2025 Internal Grants Program of Community Service (No. SPP/16/UN 63.8/PM/V/2025), which enabled the implementation of this community development initiative. Special thanks are due to the dedicated elderly cadres of Gunung Anyar District for their unwavering commitment, active participation, and exceptional collaboration throughout the program.

References

1. United Nations. Demographic Profiles Probabilistic Projections Line Charts Population Pyramids [Internet]. 2021. Available from: <https://population.un.org/wpp/graphs?loc=360&type=Demographic>
2. Jarzebski MP, Elmqvist T, Gasparatos A, Fukushi K, Eckersten S, Haase D, et al. Ageing and population shrinking: implications for sustainability in the urban century. *npj Urban Sustainability*. 2021 Dec 1;1(17).
2. United Nations D of E and SAPD. World Population Prospects: The 2015 Revision, Key Findings and Advance Tables. Working Paper No. ESA/P/WP.241. New York; 2015.

3. Franceschi C, Garagnani P, Morsiani C, Conte M, Santoro A, Grignolio A, et al. The continuum of aging and age-related diseases: Common mechanisms but different rates. *Front Med (Lausanne)*. 2018;5(61):1–23.
4. Weyand CM, Goronzy JJ. Aging of the immune system: Mechanisms and therapeutic targets. In: *Annals of the American Thoracic Society*. American Thoracic Society; 2016. p. S422–8.
5. Oh SJ, Lee JK, Shin OS. Aging and the immune system: The impact of immunosenescence on viral infection, immunity and vaccine immunogenicity. *Immune Netw*. 2019 Dec 1;19(6).
6. Kementerian Kesehatan. Laporan nasional Riset Kesehatan Dasar 2018 [Internet]. 2018 [cited 2025 Jul 15]. Available from: <https://repository.badankebijakan.kemkes.go.id/id/eprint/3514/1/Laporan%20Riset%20Kesehatan%202018%20Nasional.pdf>
7. Derajat AM, Pratidina E, Indarna AA. Hubungan pengetahuan lansia tentang diabetes mellitus dan ketidakstabilan kadar glukosa darah di Puskesmas Cibiru Kota Bandung. *Journal of Public Health Concerns*. 2025 Jan 5;4(5):154–62.
8. Rianto B, Harun O. Pengetahuan lansia tentang diabetes melitus di panti werda Cimahi berdasarkan pendidikan, umur, dan gender. *Jurnal Kesehatan Budi Luhur*. 2024;17(1).
9. Sya'diyah H, Widayanti M, Kertapati Y, Anggoro SD, Ismail A, Atik T, et al. Penyuluhan kesehatan diabetes melitus penatalaksanaan dan aplikasi senam kaki pada lansia di wilayah pesisir Surabaya. *Jurnal Pengabdian Kesehatan*. 2020;3(1):10–27.
10. Hood KK, Hilliard M, Piatt G, Ievers-Landis CE. Effective strategies for encouraging behavior change in people with diabetes. *HHS Public Access. Diabetes Manag (Lond)*. 2015;5(6):499–510.
11. Ernawati U, Andri Wihastuti T, Wiji Utami Y. Effectiveness of diabetes self-management education (DSME) in type 2 diabetes mellitus (T2DM) patients: Systematic literature review. *J Public Health Res*. 2021;10:2240.
12. Nxedlana O, Douglas M, Manu E. Strengthening community actions to improve diabetes mellitus care optimising public health facilitators. *BMC Health Serv Res*. 2025 Dec 1;25(1).
13. Christiana W, Anggraini N, Daeli NE, Kesehatan FI. Peran edukasi terhadap pengetahuan tentang manfaat labu siam pada penderita hipertensi. *E-Indonesian Journal of Health and Medical*. 2021;1:2774–5244.
14. Jaafar N, Perialathan K, Zulkepli MZ, Mohd Zin Z, Jono PE, Johari MZ. Patients' perception towards health education services received at the enhanced primary healthcare facilities: A qualitative exploration. *J Prim Care Community Health*. 2020;11:2150132720980629.
15. Suryani D, Rizal A, Eliana, Darwis, Pratiwi BA, Angraini W, et al. The effect of counseling in efforts to prevent and control non-communicable diseases. *Kemas*. 2019;14(3):297–302.
16. Buyu Prakoso A, Irawan AA, Meliasari F, Dwi Ariani E. Pencegahan kejadian risiko diabetes melitus pada lansia melalui upaya screening dan penyuluhan pengetahuan umum diabetes melitus bersama Posyandu Mardirahayu III. *Jurnal Pengabdian dan Pemberdayaan Masyarakat*. 2024;2(1):8–15.
17. Bariya B, Solanki P, Mahyavanshi D, Tolani J. Comparing the effectiveness of different health education approaches for preventing tobacco use among school-going students in a government school of Dadra and Nagar Haveli district. *J Educ Health Promot*. 2024 Jan 1;13(1).
18. Sujarwoto, Maharani A. Participation in community-based healthcare interventions and non-communicable diseases early detection of general population in Indonesia. *SSM Popul Health*. 2022 Sep 1;19:101236.
19. Susanti H, Brooks H, Yulia I, Windarwati HD, Yuliasuti E, Hasniah H, et al. An exploration of the Indonesian lay mental health workers' (cadres) experiences in performing their roles in community mental health services: a qualitative study. *Int J Ment Health Syst*. 2024 Dec 1;18(1).
20. Fang Z. The message framing effects in the process of health behavior The message framing effects in the process of health behavior change change [Internet]. 2025. Available from: https://ink.library.smu.edu.sg/etd_coll/701
21. Duckworth AL, Gross JJ. Behavior change. *Organ Behav Hum Decis Process*. 2020 Nov 1;161:39–49.
22. Kusumo MP, Kusumawati W. Barriers to understanding health education in patients with diabetes mellitus (DM) in Yogyakarta: Qualitative study. *Unnes Journal of Public Health*. 2022 Jan 31;11(1):14–22.